

Durham Research Online

Deposited in DRO:

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

van Kleunen, M. and Bossdorf, O. and Dawson, W. (2018) 'The ecology and evolution of alien plants.', Annual Review of Ecology, Evolution and Systematics, 49 (1). pp. 25-47.

Further information on publisher's website:

<https://doi.org/10.1146/annurev-ecolsys-110617-062654>

Publisher's copyright statement:

Posted with permission from the Annual Review of Ecology, Evolution and Systematics, Volume 49 © 2018 by Annual Reviews, <http://www.annualreviews.org>

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

The Ecology and Evolution of Alien Plants

Mark van Kleunen^{1,2}, Oliver Bossdorf³ and Wayne Dawson⁴

¹Zhejiang Provincial Key Laboratory of Plant Evolutionary Ecology and Conservation,
Taizhou University, Taizhou 318000, China; email: mark.vankleunen@uni-konstanz.de

²Ecology, Department of Biology, University of Konstanz, Universitätsstrasse 10, D-78464
Konstanz, Germany

³Plant Evolutionary Ecology, Institute of Evolution & Ecology, University of Tübingen, Auf
der Morgenstelle 5, D-72076 Tübingen, Germany; email: oliver.bossdorf@uni-tuebingen.de

⁴Department of Biosciences, Durham University, South Road, Durham, DH1 2LF, United
Kingdom; email: wayne.dawson@durham.ac.uk

Running title: Ecology and Evolution of Alien Plants

Corresponding author: Mark van Kleunen, mark.vankleunen@uni-konstanz.de

Keywords

biotic interactions, exotic plant, invasive, macro-ecology, naturalized, rapid evolution

Abstract

We review the state-of-the-art of alien plant research with emphasis on conceptual advances, and knowledge gains on general patterns and drivers, biotic interactions and evolution. Major advances include the identification of different invasion stages and invasiveness dimensions ([geographic range, habitat specificity, local abundance](#)), and the need for appropriate comparators while accounting for [propagule pressure and ~~introduction history~~ year of introduction](#). Developments in phylogenetic and functional-trait research, ~~and hybrid modelling~~, bear great promise [for better understanding of the underlying mechanisms](#). Global patterns are emerging with propagule pressure, disturbance, increased resource availability and climate matching as major invasion drivers, but species characteristics also play a role. Biotic interactions with resident communities shape invasion outcomes, with major roles for species diversity, enemies, novel weapons and mutualists. There is mounting evidence for rapid evolution of invasive aliens and evolutionary responses of natives, but a mechanistic understanding will require better integration of molecular and phenotypic approaches. We hope the open questions identified will stimulate further research on the ecology and evolution of alien plants.

1. INTRODUCTION

The study of alien organisms and their biotic interactions and varying invasion success is a major research area in ecology and evolutionary biology. Its motivation has always been two-fold: On the one hand, scientists and conservation managers have been concerned about negative impacts of alien organisms on native biodiversity and economy. On the other hand, since alien organisms often experience novel ecological contexts, and there is large variation in invasion success, which is at least partly explained by ecological and evolutionary processes, the study of alien species greatly advances our fundamental ecological and evolutionary understanding (Sax et al. 2007). The initial research agenda for invasion biology was set by two seminal books on the ecology (Elton 1958) and genetics (Baker & Stebbins 1965) of invasive species. Research has grown exponentially particularly in the second half of the previous century (Gurevitch et al. 2011), and invasion biology is now a mature discipline.

Within invasion biology, the study of alien plants has been particularly strong, with its findings summarized in numerous reviews (e.g., Rejmánek 1996, Pyšek & Richardson 2007). Nevertheless, our understanding of alien plant invasions, and invasion biology more broadly, has long been hampered by unclear and inconsistent use of definitions (Pyšek et al. 2004), [failure to account for year of introduction and propagule pressure \(and a lack of appropriate null models](#) (Colautti et al. 2006), [and use of comparator groups](#) [comparisons of invasive species \(or populations\) to reference species \(or populations\) that do not address the research question](#) (van Kleunen et al. 2010a). Moreover, there has sometimes been a lack of understanding of how different hypotheses in plant invasion biology are related (Catford et al. 2009). In recent years, there has been much progress in this regard.

In this review, we describe some of the major conceptual and methodological advances, and empirical studies that have improved our understanding of plant invasions. We

do not provide a systematic, exhaustive review, but ~~a series of~~ rather an overview of selected key topics where progress has been made, from macro-ecology and biotic interactions to evolution and genetics. Our review is accompanied by a visual summary in **Figure 1** where we indicate, for each topic, how well it has been studied so far, how consistent the results were and how many open questions there still are in it have been from study to study, and how many questions remain open. Although this figure is clearly somewhat subjective, we hope that together with the ‘way-forward’ sections below, it will stimulate and guide future research on the ecology and evolution of alien plants.

2. CONCEPTUAL AND METHODOLOGICAL ADVANCES

With the rapid increase in research on alien plants, our understanding of the processes that drive plant invasions has increased considerably. Several conceptual and methodological advances have contributed to this.

2.1. The Invasion-stages Framework

Invasion biology has developed a large vocabulary with multiple terms for the same things (e.g., alien, exotic, non-native, non-indigenous), and multiple definitions for the same terms (e.g., invasive; see below). To increase clarity and consistent use of terminology, Richardson et al. (2000) proposed an invasion-stages framework with a sequence of barriers that a plant species has to overcome to become invasive (**Figure 2**). In this framework, aliens are plant species that have passed a biogeographic barrier (e.g., an ocean) with help of humans. The aliens that have passed the subsequent environmental barrier and sometimes occur in the wild but do not form persistent populations are considered casuals. The ones that have passed the environmental and reproductive barriers, and have established wild populations that persist

Formatted: Font: Bold

over multiple life cycles, are considered naturalized. The [subset of](#) naturalized species that have overcome the dispersal barrier [and produce reproductive offspring, often in very large numbers, at considerable distance from parent plants \(>100 m within <50 years; Richardson et al. 2000\)](#) ~~within the non-native range and subsequent environmental barriers they encountered~~ are considered invasive.

While the Richardson et al. framework is widely applied (3042 citations in Google Scholar, accessed 11 January 2018), definitions of ‘invasive’ vary. The Richardson et al. (2000) definition is neutral with respect to ecological and economic impacts, whereas the Convention on Biological Diversity (CBD 2000) defines invasive species as those whose introduction and/or spread threatens biological diversity. Problematic is that the latter is frequently unknown. Other definitions of ‘invasive’ are used less frequently, although they consider interesting aspects. For example, ~~Alpert et al.~~ [\(2000\)](#) proposed decoupling ‘invasive’ from being alien, and Hufbauer & Torchin (2007) proposed defining a species invasive when its demographic performance is higher in the invaded than in the native range. While many alien species might qualify as invasive under all these definitions, there are exceptions. Parker et al. (2013) showed that plants among ‘*100 of the world’s worst invasive alien species*’, which follows the CBD (2000) definition, perform on average better in their non-native than in their native ranges, thus meeting the Hufbauer & Torchin (2007) ‘invasive’ definition. However, there was high variability among these species in this regard. Although none of the ‘invasive’ definitions is necessarily better than the others, the use of multiple definitions has resulted in confusion, and might explain some of the apparently conflicting findings among studies. Therefore, we call for researchers to always state clearly which definition of ‘invasive’ is used. Here, we use the Richardson et al. (2000) definitions of alien, naturalized and invasive, [unless stated otherwise](#).

2.2. The Multiple Dimensions of Invasiveness

The multiple 'invasive' definitions demonstrate that invasive species, as described in the literature, are not a homogeneous group. While the Richardson et al. (2000) definition of invasive primarily focuses on spread of the naturalized species, it also implicitly ~~also~~ assumes that the species are locally abundant (i.e., produce reproductive offspring in large numbers). Species show continuous variation in spread (i.e., range size) and abundance, and which means that invasiveness is not binary, but is instead ~~therefore a continuous variable~~ and multidimensional, ~~rather than a binary (yes/no) variable. Moreover, w~~ While some naturalized alien species are wide-spread, they have a low locally abundant abundance; they might have a small non-native range or be restricted to few habitats, whereas others might have a large-small range and occur in many habitats but have sparse populations have a high local abundance. ~~In other words, invasiveness has multiple dimensions (Figure 2). For~~ the related concept of ~~The idea that rarity and commonness (or rarity), Rabinowitz (1981) proposed, have three dimensions — in addition to geographical range, and habitat specificity,~~ local population size (i.e., abundance), to also consider the range of habitats in which a species occurs (i.e., habitat generality). This idea of multiple dimensions of a species' distribution or commonness ~~— was originally developed by Rabinowitz (1981), but~~ has only recently been applied to invasiveness of alien plants (Dawson et al. 2013, Catford et al. 2016).

Formatted: Font: Bold

The different invasiveness dimensions are likely to be associated with different main barriers (Figure 2). Range size (spread) may be mainly determined by one or more dispersal barriers, local abundance by one or more competition barriers, and habitat generality by one or more environmental barriers. Depending on the specific research question, one could also consider ~~Potential additional further~~ dimensions, ~~such as -of invasiveness are the rate of~~ spread ~~rate~~ (Catford et al. 2016), and different categories of ecological and economic impacts,

Formatted: Indent: First line: 1.27 cm

Formatted: Font: Bold

which would then also include the CBD (2000) definition of invasive. We believe that consideration of these different invasiveness dimensions in research on alien plants may provide more consistent results among studies prevent researchers from mistakenly concluding that results are in conflict.

2.3. Comparative Approaches for Assessing Determinants of Invasion Success

The variety in approaches used to study alien plants and their invasion success is enormous. While there are many descriptive studies on individual invasive species, the question of what determines invasion success requires a comparative approach. Such studies are most powerful when they include many species, populations and sites (van Kleunen et al. 2014). The comparator choice is pivotal to whether the question of interest can be answered (van Kleunen et al. 2010a). Although most studies on traits associated with invasiveness have compared invasive alien to native species (Pyšek & Richardson 2007, van Kleunen et al. 2010b, Davidson et al. 2011), this cannot reveal why alien species differ in invasiveness. This requires a comparison between alien species that differ in their degree of invasiveness between invasive and non-invasive alien species, either along a single dimension or along multiple dimensions of the invasiveness continuum (Figure 2). When these comparisons include non-invasive alien species, though it is also important to realize that the latter may be at different invasion stages. Some non-invasive aliens, some might not be not found outside of cultivation, some are just casuals and others are naturalized but not invasive have been introduced, and others might be introduced but not naturalized (van Kleunen et al. 2015b Figure 2). As different traits might be associated with each of these stages (Dietz & Edwards 2006, Dawson et al. 2009, van Kleunen et al. 2015b), the results of the comparison between invasive and non-invasive aliens may strongly depend on the invasion stage of the latter.

Formatted: Font: Bold

Formatted: Font: Bold

125

126 **2.4. Introduction History as a Null Model**
127 Accounting for propagule pressure and year of introduction

128 It seems almost trivial that alien species introduced in greater numbers or more frequently are
129 more likely to naturalise and become invasive, and thus should be accounted for.
130 Nevertheless, the need to account for this so-called propagule pressure has been ~~formalized as~~
131 a ‘null model’ for invasion success recognized only recently (Colautti et al. 2006). Similarly,
132 alien species that were introduced earlier should have had more opportunities to naturalize
133 and become invasive (Rejmánek 2000). We will discuss the importance of propagule pressure
134 and year of introduction in more detail below. However, accounting for ~~the introduction~~
135 history propagule pressure and year of introduction of ~~an~~ alien species has been important for
136 disentangling the ecological and evolutionary processes that contribute to plant invasions.

137

138 **2.5. Darwin’s Naturalization Conundrum, Scale Dependency and Coexistence Theory**

139 There has been long-standing interest in how differences between alien and native plants
140 determine invasion success. Darwin (1859) hypothesised that alien plants distantly related
141 from the native communities are more likely to naturalize. A mechanism underlying
142 Darwin’s naturalization hypothesis (Rejmánek 1996) could be stronger niche differentiation
143 between resident natives and more distantly related aliens (Thuiller et al. 2010). In addition,
144 the more distantly related the alien plant is, the less likely it is that herbivores and pathogens
145 will spill over from native residents (see Enemy Release section below). Darwin (1859) also
146 hypothesised that alien species from genera that occur in native regional floras may be more
147 likely to naturalize because they share ~~the~~ the same pre-adaptations as the related natives.
148 These seemingly contradictory hypotheses are now referred to as ‘Darwin’s naturalization
149 conundrum’ (Thuiller et al. 2010).

The realization that different ecological processes act at different spatial scales ~~is~~
~~helping~~ might help to resolve this conundrum ~~and as well as the apparent discrepancies~~
~~between studies that found positive, negative or no significant relationships between~~
~~naturalization success of alien plants and their the mixed findings~~ phylogenetic relatedness to
~~native plants~~ (Thuiller et al. 2010). Whereas abiotic environmental filtering, which requires
pre-adaptation, acts at all spatial scales, biotic filtering acts only at the small scales where
species interact (Thuiller et al. 2010, Gallien & Carboni 2016). Therefore, Darwin's
naturalization hypothesis (i.e., the hypothesis that alien species distantly related to native
species are more likely to naturalize) should only operate at small spatial scales. As the
different processes might act simultaneously, the relationship between invasion success and
phylogenetic distance might actually be non-linear and depend on the phylogenetic scale
(Thuiller et al. 2010, Gallien & Carboni 2016). These ideas, however, still need to be tested
explicitly.

Another potential reason for the mixed findings is that the expected relationship
between phylogenetic distance and invasion success rests on the assumption that
phylogenetic distance reflects niche differentiation due to trait differences. However, not all
traits are phylogenetically conserved. Moreover, once a species has passed the abiotic
environmental filter, the outcome of competition is not only determined by niche differences
but also by fitness differences (Chesson 2000, Mayfield & Levine 2010). The latter also
depend on trait differences (Mayfield & Levine 2010). This modern coexistence theory has
only recently been extended to biological invasions and linked to Darwin's Naturalization
naturalization ~~Conundrum~~ ~~conundrum~~ (MacDougall et al. 2009, Thuiller et al. 2010). While
empirical studies are still rare, the recent advances in theory, tools and data for phylogenetic
and functional trait analysis bear great promise to better understand invasions into plant
communities.

175

176 **2.6. Modelling Plant Invasions**

177 ~~Once an alien species has been introduced in sufficient numbers, its invasion success is~~
178 ~~contingent upon~~Studies on invasion success usually focus either on invasibility of the
179 environment ~~and or~~ invasiveness of the species (Richardson & Pyšek 2006). In recent
180 decades, more data has become available on regional differences in numbers of species
181 invading, and species differences in invasion success. Together with data on environmental
182 and socio-economic variables, species characteristics and introduction history, this
183 information has revealed some general patterns providing insights into processes underlying
184 plant invasions.

185

186 **3.1. A Global Overview of the Naturalized Alien Flora**

187 A recent compilation of >800 regional naturalized alien floras revealed that globally >13,000
188 plant species have become naturalized (van Kleunen et al. 2015a, Pyšek et al. 2017). In other
189 words, ~4% of the extant vascular flora has expanded beyond species' native ranges with
190 help of humans. Most of these naturalizations occurred in the last two centuries, and although
191 the rate at which newly naturalized alien species are discovered has plateaued, it is not
192 decreasing yet (Seebens et al. 2017). We can thus expect many more new naturalizations in
193 the next decades, ~~and a major research question is, how will they will be distributed around~~
194 ~~the globe is one of the main questions (Figure 1)?.~~

Formatted: Font: Bold

195 North America has the highest number of naturalized plants, Antarctica has the lowest
196 number, and the highest density occurs on the Pacific islands (van Kleunen et al. 2015a). The
197 Northern Hemisphere continents are the major donors of naturalized species. Europe has even
198 donated 288% more species than would be expected considering its small native flora (van

199 Kleunen et al. 2015a), probably as a consequence of European colonialism. Hotspots of plant
200 invasions are found on islands and in coastal regions (Dawson et al. 2017, Pyšek et al. 2017).
201 For mainland regions, naturalized plant species richness increases with decreasing mean
202 annual temperature, and with increasing mean annual precipitation, human population density
203 and per capita gross domestic product (Dawson et al. 2017). The importance of economic
204 factors, specifically past bilateral trade, leads to the forecast that emerging economies, such
205 as India and China, will see a rise in plant naturalizations in the next two decades (Seebens et
206 al. 2015), [and are also likely to become more important donors of naturalized species](#).

207 For invasive plants *sensu* Richardson et al. (2000)~~(i.e., rapidly spreading alien plants;~~
208 [see section 2.1](#)), few data on global patterns are currently available. A global analysis of
209 invasive trees and shrubs showed, as for naturalized plants in general, that the highest
210 numbers are found in North America and the Pacific Islands (Rejmánek & Richardson 2013).
211 For invasive plants *sensu* CBD (2000)~~(i.e., alien species threatening native biodiversity; see~~
212 [section 2.1](#)), Pyšek et al. (2017) similarly showed that there is a strong correlation between
213 the numbers of invasive and naturalized species in a region. Nevertheless, more data on
214 invasive species *sensu* Richardson et al. (2000) [and the different dimensions of invasiveness](#)
215 [\(Figure 2\)](#) are needed to better understand the global patterns and drivers of plant
216 invasiveness.

Formatted: Font: Bold

218 3.2. Propagule Pressure

219 Propagule pressure – a composite measure of the number of release events and the number of
220 individuals released per event – is considered the most consistent driver of invasion success
221 (Lockwood et al. 2005, 2007, Simberloff 2009). Theoretically, a high propagule pressure
222 increases the likelihood of overcoming Allee effects, and demographic and environmental
223 stochasticity (e.g., Shea & Possingham 2000). Empirical studies on propagule pressure of

224 alien plants mostly use proxies such as sales or planting frequencies of ornamental plants
225 (Dehnen-Schmutz et al. 2007, Feng et al. 2016, Maurel et al. 2016), and are correlative in
226 nature. Introduction experiments, on the other hand, have the disadvantage that ~~they last~~
227 [duration is](#) too short to capture the full invasion process. Nevertheless, the few available
228 introduction experiments confirm that propagule pressure has a strong positive effect on early
229 establishment of alien plants (e.g., Von Holle & Simberloff 2005, Kempel et al. 2013).

230 Despite the importance of propagule pressure, little is known about the roles of its
231 elements. In theory, many released individuals per introduction event should help
232 overcoming demographic stochasticity and Allee effects, whereas many introduction events
233 should facilitate overcoming environmental stochasticity (Shea & Possingham 2000,
234 Simberloff 2009). Experimental studies separating these propagule-pressure elements are rare
235 and restricted to animals (e.g., Sinclair & Arnott 2016). Genetic variation, [which is likely to](#)
236 [increase with each additionally introduced individual](#), is another element of propagule
237 pressure, and has been shown to play a role in *Spartina alterniflora* invasion in China (Wang
238 et al. 2012). More experimental studies are needed that consider the roles of propagule
239 pressure's different elements in alien plant invasions ([Figure 1](#)), and also how they interact
240 with other drivers.

241

242 3.3. Human disturbance

243 Disturbance [by humans](#) is thought to be another major driver of plant invasions (Lockwood et
244 al. 2007). Disturbance is defined as any relatively discrete event in time that disrupts
245 ecosystem, community or population structure, and changes resources, substrate availability
246 or the physical environment (White & Pickett 1985). Many disturbances are naturally
247 recurring events, and a change in disturbance regime [by humans](#) rather than the disturbance
248 event itself may promote invasions (Hobbs & Huenneke 1992). This complexity makes

249 generalizing across disturbance types difficult (Lockwood et al. 2007). Nevertheless, global
250 (Dawson et al. 2017) and regional (e.g., McKinney 2001) analyses show that richness of
251 naturalized and invasive plants are usually strongly associated with human-population
252 density, which is ~~arguably a good surrogate for~~associated with human disturbance.

253 In a review, Colautti et al. (2006) showed that the majority of plant studies found that
254 disturbance, usually involving the destruction of biomass, increases invasibility of plant
255 communities. Most studies considered only invaders already in the system, which can
256 obviously invade (Lockwood et al. 2007), and manipulated disturbances natural to the
257 system. Recent experiments in which locally non-occurring species were introduced to native
258 grassland communities showed that establishment from seed is considerably higher after
259 tilling of the soil (i.e., a human disturbance; e.g., Kempel et al. 2013). Overall, there is thus
260 good evidence that human disturbances promote biological invasions. However, which alien
261 species benefit the most from disturbance and why, remains an open question (Figure 1).

Formatted: Font: Bold

262 3.4. Responses to Additional Resources

263 Many disturbance events and anthropogenic global change drivers ~~may~~ change resource
264 availabilities. As predicted by the fluctuating-resource-availability hypothesis (Davis et al.
265 2000), increases in resource availability make plant communities more susceptible to plant
266 invasion (Seabloom et al. 2015). However, not all alien plants take advantage of increased
267 resources; so successful alien plants may be those that capitalize most strongly on increased
268 resources. ~~(Davidson et al. 2011) showed in a meta-analysis that invasive species were more~~
269 ~~plastic in growth, morphology and physiology than native species, but this did not result in~~
270 ~~fitness advantages. However, a~~ multi-species experiment showed that among native and
271 among alien species in Switzerland, common species capitalized more on nutrient increases
272 than rare species (Dawson et al. 2012a). Similarly, a meta-analysis showed that globally more
273 widespread alien species exhibited greater biomass responses to increases in resources

274 (Dawson et al. 2012b). So, although some alien plants invade low-resource environments
275 (Funk 2013), many invasive alien plants have a high capacity to capitalize on increased
276 resource levels.

277 Environmental change is frequently characterized by a change in variability as well as
278 in mean conditions. For example, floods and fires may temporarily increase nutrient
279 availability. Parepa et al. (2013) showed that invasive *Fallopia* spp. grown with native plants
280 benefited more strongly when supplied with nutrient pulses instead of a constant supply. In a
281 multi-species experiment, naturalized alien plants produced more biomass when nutrients
282 were provided as a single large pulse in the middle of the growth period compared to plants
283 with a constant supply (Liu & van Kleunen 2017). The reverse was true for native plants.
284 Thus, many successful alien plants are opportunists that take advantage of nutrient pulses.

285 The underlying mechanisms still need more research (Figure 1).

Formatted: Font: Bold

287 **3.5. Lag Phases and Invasion Debts**

288 Species need time to move from one invasion stage to the next. Once an alien plant has been
289 introduced, there is a lag phase before it ~~becomes~~ ~~naturalizes~~ ~~naturalized~~, and one before it ~~is~~
290 ~~considered to be~~ ~~becomes~~ invasive (~~i.e., starts to accelerate its spread~~) (Crooks 2005).

291 Unfortunately, few studies distinguish between the two lag phases, and no study quantified
292 both. Lag phases may simply be an inherent characteristic of exponential population growth,
293 or they may result from Allee effects or time needed for evolutionary adaptation or
294 environmental change (Crooks 2005). A lack of hard data prevents us from understanding the
295 importance of these mechanisms.

296 For ornamental and forestry species, the introduction-naturalization lag phase ranges
297 from two to over 370 years (Kowarik 1995, Binggeli 2000, Caley et al. 2008, Daehler 2009).
298 The few studies that quantified this lag phase indicate that it is shorter in tropical (Binggeli

299 2000, Daehler 2009) than in temperate (Kowarik 1995, Caley et al. 2008) regions, possibly
300 because of year-round growth in the tropics (Daehler 2009). These studies further clearly
301 show that shorter generation times result in shorter lag phases, suggesting that differences in
302 lag phases may diminish if measured in terms of generation times.

303 The naturalization-invasion lag phase has been quantified using time series of
304 herbarium records. Most of these lag phases are <50 years (Aikio et al. 2010, Larkin 2012).
305 However, these lag phase estimates should be interpreted with caution as Hyndman et al.
306 (2015) argued that [several](#) assumptions underlying the lag-phase-estimation approach were
307 violated. [For example, because the lag-phase estimation involves fitting models to cumulative](#)
308 [numbers of herbarium records over time, the assumption of non-independence of the](#)
309 [residuals is violated.](#) Future studies on naturalization-invasion lag phases should aim to
310 optimize statistical approaches, and then assess which factors drive variation in lag phases
311 [\(Figure 1\).](#)

Formatted: Font: Bold

312 The existence of lag phases implies that the current numbers of naturalized and
313 invasive alien species are determined by processes in the past. Indeed, Essl et al. (2011)
314 demonstrated that current plant naturalization patterns in Europe are better explained by
315 socioeconomic factors from the year 1900 than from the year 2000. Research on such so-
316 called invasion debts (Seabloom et al. 2006) will thus be useful for predicting future
317 invasions.

318

319 **3.6. Climatic Suitability**

320 Plant distributions have long been thought to be primarily restricted by biogeographic barriers
321 and climatic conditions (Good 1931). Consequently, climatic suitability is considered a major
322 predictor of plant invasion success (Panetta & Mitchell 1991). However, recently, several
323 studies reported that ~40% (Guisan et al. 2014) or even >65% (Atwater et al. 2018) of

invasive species have undergone climatic niche shifts. The reasons for these shifts require more research (Figure 1), but they call into question the assumption of climatic niche conservatism ~~into question~~. This suggests that estimates of climatic suitability from the native range may be poor predictors for invasion success. Surprisingly, few studies have explicitly related the modelled climatic suitability of introduced alien plants to their actual invasion success. The naturalization success of 449 Chinese woody species introduced to Europe was significantly correlated with climatic suitability (Feng et al. 2016). However, in that study, climatic suitability only explained 5% of the variance in naturalization success, possibly due to the large geographical extent (Europe) considered. On the other hand, a study on the garden flora of a small municipality in southern Germany found that climatic suitability was very strongly related to local naturalization success (Mayer et al. 2017). So, overall climatic suitability seems to be an important prerequisite for plant invasions.

Formatted: Font: Bold

3.7. Species Characteristics

Baker (1965)'s list of 'ideal weed' characteristics was the starting point for research on species characteristics related to invasion success. Several reviews (Pyšek & Richardson 2007, van Kleunen et al. 2015b) and meta-analyses (van Kleunen et al. 2010b, Davidson et al. 2011) have summarized the results. Although some trends appear, results depend on whether invasive aliens are compared to natives or non-invasive aliens (van Kleunen et al. 2010b), and on the invasion stage considered (Dietz & Edwards 2006, Dawson et al. 2009). Nevertheless, a few characteristics are globally associated with naturalization success. Using a global database on breeding systems of 1752 plant species, Razanajatovo et al. (2016) showed that species with an increase in greater self-fertilization ability, the number of were naturalized in more regions around the world globally in which a species is naturalized increased. Furthermore, species listed in databases as harmful invasive species (i.e.,

following the CBD (2000) definition of invasive; see section 2.1) are characterized by ~~Small~~ a small genome size and polyploidy are plant characteristics highly associated with invasion success (Pardoll et al. 2014). As genome size and polyploidy are functional traits, we ~~will~~ need approaches such as path analysis to separate direct and indirect effects of functional traits associated with genomic characteristics on invasion success.

The general lack of consistent relationships between species characteristics and invasion success limits predictions of invasions based on those characteristics. This does not mean that species characteristics are unimportant. Predictive power may improve by considering invasion stage, ~~invasiveness dimension~~, spatial scale and environmental context (van Kleunen et al. 2015b), by testing whether different species traits are associated with the different dimensions of invasiveness (Figure 1; Catford et al. 2016). Studies also need to test for interactions between traits, and for non-linear relationships between invasion success and traits. Moreover, they should account for introduction bias, i.e., the phenomenon that species with certain characteristics were introduced earlier or more frequently, and consequently have higher apparent invasion success (Lockwood et al. 2005, Colautti et al. 2006, Maurel et al. 2016).

Formatted: Font: Bold

3.8. The Way Forward for Research on General Patterns and Drivers of Invasion

Success

Recently built databases describing the global distribution of naturalized and invasive alien plants allow researchers to address major macro-ecological questions on invasion success of alien plants. The geographical units for these databases are administrative regions of different sizes. Ideally, such data should include the habitat-affiliations within the regions, or should be collected for grid-cells/geo-referenced units of equal size. Another step forward would be to integrate the distributional data of naturalized alien and native species. This would allow ~~assessing~~ assessment of the importance of habitat filtering, and how much invasions

374 contribute to homogenization of the global flora. Furthermore, to better understand the
 375 importance of species characteristics and historical factors in invasions, we need data on alien
 376 species that were introduced but failed to ~~establish~~ [naturalized \(Dehnen-Schmutz et al. 2007,](#)
 377 [Maurel et al. 2016\)](#), at least until now. These data will also allow better quantification of
 378 invasion debt. Compilations of nursery catalogues and botanical garden inventories [from](#)
 379 [around the world](#) will be a good starting point for a global introduced alien flora.

380 A major objective of invasion biology is to understand the mechanisms driving
 381 invasion success, but studies on global patterns are correlative, not causative. Thus, some of
 382 the identified apparent drivers of plant invasion patterns might not be true drivers. For
 383 example, because propagule pressure is a major driver of invasion success, it should be ~~used~~
 384 ~~as a null model~~ [accounted for](#) when testing the roles of e.g. species characteristics. However,
 385 it could be that the propagule pressure only appears to be important because species with
 386 characteristics that promote invasion success have been introduced more frequently (Maurel
 387 et al. 2016). Structural equation modelling (Shipley 2000) offers a [potential](#) statistical
 388 solution to better separate the direct and indirect drivers of plant invasions. Causative tests for
 389 generality of invasion drivers could further be achieved by establishing research networks
 390 ~~that require only~~ [in the Native North America and Europe \(Boedeltje et al. 2014\)](#)

392 4. BIOTIC INTERACTIONS OF ALIEN PLANTS

393 Alien plants interact with native resident plants and other organisms. The resulting effects on
 394 alien plant performance and fitness determine whether a species is able to establish in a local
 395 community (Levine et al. 2004, MacDougall et al. 2009). Biotic interactions occur at the
 396 individual plant scale, but should affect invasion success at larger scales. For instance, altered
 397 biotic interactions in the introduced compared to the native range may modulate the realised

398 niches of invasive plants in the introduced range, possibly leading to habitat expansion or
399 climatic niche shifts (Guisan et al. 2014, Atwater et al. 2018).

400

401 **4.1. Diversity of Resident Communities**

402 Areas with high native species richness frequently also have high alien richness (Stohlgren et
403 al. 2003). This likely results from habitat heterogeneity covarying with both native richness
404 and alien richness at landscape scales (Levine et al. 2004). In contrast, at local scales – as
405 predicted by Elton (1958)’s diversity-invasibility hypothesis – more species-rich
406 communities are frequently more resistant to invasions (Levine et al. 2004). Most likely, this
407 is because a more species-rich community occupies more of the available niche space.

408 Although even the most diverse communities cannot resist invasions completely, there is
409 strong support for the diversity-invasibility hypothesis (Levine et al. 2004). However, we still
410 require studies involving communities not dominated by herbaceous species [to assess the](#)
411 [generality of the pattern](#), and studies testing the importance of phylogenetic and functional
412 diversity [to better understand the mechanisms underlying the pattern \(Figure 1\).](#)

413

414 **4.2. Enemy Release**

415 Alien plants may be released from herbivores and pathogens, especially from specialists [that](#)
416 [have not been co-introduced](#), resulting in a competitive advantage over natives. This so-called
417 enemy-release hypothesis (Elton 1958, Keane & Crawley 2002) is perhaps the best known
418 hypothesis in invasion ecology. Indeed, some invasive alien plants have fewer enemies
419 associated with them and incur less damage in the introduced than in the native range (e.g.,
420 Mitchell & Power 2003, Meijer et al. 2016). However, results from studies comparing enemy

Formatted: Font: Bold

421 damage on alien and co-occurring native species are equivocal, and many studies do not
422 ~~link assess how~~ damage affects fitness of the to alien plant ~~performances~~ (Chun et al. 2010).

423 Most enemy-release studies focus on aboveground enemies, but the role of soil
424 pathogens has gained attention (Dawson & Schrama 2016). Some European native plants
425 invasive in North America grow better in sterilized than in non-sterilized soils, but only for in
426 European soils, and but not in North American soils, ~~after soil sterilisation~~ (Maron et al.
427 2014). This suggests that the European plants do not suffer from soil pathogens in North
428 America. Such 'plant-soil-feedback' studies, however, treat soil as a black box. One
429 exceptional example is a study on *Prunus serotina*, which in its native North American range
430 suffers from high seedling mortality due to infection by specific oomycete pathogens, while
431 in Europe these are less virulent (Reinhart et al. 2010). We need more such studies that verify
432 which microbes have negative effects on plants and are absent in the introduced range.

433 Enemy release might decrease with time as enemies in the invaded range may shift
434 hosts and start to utilise the invader (Diez et al. 2010). ~~There are reports, however, that some~~
435 ~~generalist soil pathogens that increased in abundance under invasive plants, spill over to~~
436 ~~infect native plants (e.g., *Chromolaena* in India; Mangla et al. 2008). Therefore, the outcome of these novel encounters with~~
437 However, new encounters at the expanding edges of invasions may also result in temporary
438 enemy release, if the natural enemies there do not utilise the invader immediately. So,
439 whether alien plants escape or recruit enemies at the range edges remains an open question
440 (Fig. 1). Continued enemy release may occur if the natural enemies do not utilise the invader
441 example, generalist soil pathogens that can't increased in abundance under invasive plants, spill
442 over to infect native plants and thereby enhance invader dominance (e.g., *Chromolaena*
443 *odorata* in India; Mangla et al. 2008). Therefore, the outcome of these novel encounters with
444 enemies is unknown hard difficult to predict for many alien plants.

Commented [WD1]: I think elsewhere we use 'alien' rather than 'introduced', but either way, I guess we should be consistent.

Formatted: Font: Bold

4.3. Novel Weapons

The novel weapons hypothesis posits that some chemical compounds of an alien species can negatively affect naïve native species (Callaway & Ridenour 2004). These novel weapons may be released as root exudates, or as leachates from living or dead plant material. The most famous example is the European plant *Centaurea stoebe*, which has allelopathic effects on native plants in the North American grasslands it invades, but not on competing plants in its native European range (Thorpe et al. 2009). The allelopathic effect of *C. stoebe* has been demonstrated in the field (Thorpe et al. 2009), but such field tests of the novel weapons hypothesis are rare.

Novel biochemistry may also have impacts on other organisms (Cappuccino & Arnason 2006). For example, novel defence chemicals may reduce herbivory by naïve herbivores (Macel et al. 2014), but there is no evidence that invasive alien plant are generally more deterrent to generalist herbivores than native plants are (Lind & Parker 2010). Some novel chemicals can also suppress mycorrhizal fungi beneficial to native competitors but not required by the invader, as shown for *Alliaria petiolata* in North America (Stinson et al. 2006). Over time, ~~the novelty of these weapons may wear off~~ the native organisms may adapt to the novel chemicals (Lankau et al. 2009), but more research is needed to understand ~~ecological and evolutionary~~ the processes involved. Another question that deserves research attention is whether ~~novelty of native plant~~ chemical compounds of native plants are novel to alien plants and contribute to increases resistance against invasion (**Figure 1**; Cummings et al. 2012).

Formatted: Font: Bold

4.4. Mutualists

Mutualisms of plants with soil microbes, pollinators and seed dispersers may influence invasions, but have received less attention than enemies (Traveset & Richardson 2014).

Therefore, their relative importance remains unknown (Figure 1). Nitrogen-fixing microbes and mycorrhizal fungi are the two main groups of soil mutualists. Some invasive-alien plants have profited from are likely to have become invasive due to their ability of having nitrogen-fixing root symbionts, particularly when N-fixing is absent in the native plant-species pool (Vitousek & Walker 1989). Alien plants may acquire the N-fixing microbes through co-introduction or ‘ecological fitting’ of alien plants to native N-fixers-fixing microbes (Le Roux et al. 2017), leading to shifts in N-fixer community composition between the native and alien ranges. The latter is indicated by differences. However, it is unclear if these shifts in N-fixing rhizobial community composition between native and alien ranges of plants hinder or enhance invasion (Körner, 2017; Stoll, 2016). However, the relative importance of N-fixing microbes in the native and alien ranges of plants remains unknown. Therefore, the relative importance of N-fixing microbes in the native and alien ranges of plants remains unknown.

As most plants have mycorrhizal fungal associations that help with the uptake of nutrients (Wang & Qiu 2006), invasion success may depend on mycorrhiza. Indeed, some tree invasions in South America depended upon the co-introduction of ectomycorrhizal fungi (Hayward et al. 2015). In Germany, mycorrhizal, and particularly facultative mycorrhizal, alien plants have a wider distribution than non-mycorrhizal species (Menzel et al. 2017). It is not yet known whether this relationship holds globally.

The vast majority of flowering plant species are pollinated by animals (Ollerton et al. 2011), but surprisingly few studies have explicitly assessed the importance of plant-pollinator mutualisms for plant invasions (Stout & Tiedeken 2017). Pollen limitation is relatively uncommon among invasive plants (Pyšek et al. 2011), possibly because of high [autofertility](#) [self-fertilization ability](#) (Razanajatovo et al. 2016) and ability to integrate in native plant-pollinator networks (Vilà et al. 2009). Surprisingly, Razanajatovo & van Kleunen (2016) found that non-naturalized alien species are also not pollen-limited. More studies are needed to test whether this is a general phenomenon. Few studies of plant-pollinator interactions (e.g., Stout et al. 2006) and [selfing-self-fertilization](#) rates (e.g., Ollerton et al. 2012) have

compared reproductive success in native and alien range populations. [More studies would be needed to](#) shed light onto whether shifts in pollinators or [selfing-self-fertilization](#) in the alien range contribute to invasion success.

Alien plants are more likely to spread into (semi-)natural habitats if they recruit native fruit-consuming animals as seed dispersers (e.g., Cordeiro et al. 2004). However, the general importance of dispersal relative to other factors, and how plant-disperser dynamics change over time are still poorly understood. [In a The results of a study in rare dispersal experiment in Canada suggest that the invasive ant *Myrmica rubra*, as well as the native ant *Aphaenogaster rudis*, contribute to seed dispersal of the native *Cubibimide* species field origin in a portable microclimate \(Pied 2014\).](#) Introduced seed-dispersing animals [can](#) also facilitate invasions by alien plants in [Hawaii](#) [Hawaii](#)’, where fruits of *Myrica faya* are dispersed by the alien bird *Zosterops japonica* (Vitousek & Walker 1989). Thus currently non-naturalized plants may still pose a future invasion risk if a suitable disperser [gets is subsequently introduced, leading to ‘invasional meltdown’ \(Simberloff & Von Holle 1999\).](#)

4.5. The Way Forward for Research on Biotic Interactions of Alien Plants

Most research on biotic interactions as [drivers](#) [determinants](#) of alien plant success has focused on single interaction types, when in reality, multiple interactions occur simultaneously. There have been calls to consider multi-trophic interactions centred on alien plants (Harvey et al. 2010), though we have yet to move beyond the use of model interactors (often generalists) under greenhouse conditions (Kempel et al. 2013). While research on the role of plant-soil feedbacks in invasions is expanding, we often do not know which types of micro-organisms are the most important ‘players’ contributing to net soil-biota effects (Dawson & Schrama 2016). Progress here requires detailed studies that involve isolation, identification and re-inoculation of plants with putative soil pathogens and mutualists. We also recommend that

520 more attention be paid to the role of mutualists of all types in invasions, in order to rebalance
521 the current bias toward natural enemies.

522 Biotic interactions do not operate in a vacuum, and are contingent upon the traits and
523 evolutionary history of both the alien plant and its putative interactors in the introduced
524 range. Interactions may change over time as a result of ecological ~~or~~ and evolutionary
525 processes. Thus, to better understand how biotic interactions have affected and will continue
526 to affect invasions, we need to consider both evolutionary history, and current eco-
527 evolutionary dynamics.

528

529 **5. EVOLUTION AND GENETICS OF INVASIVE PLANTS**

530 There is considerable potential for evolution resulting from plant introductions to new ranges.

531 Alien plants ~~are less adapted to their new environments than to their native environments,~~
532 ~~they may experience novel conditions, and thus selection pressures.~~ lack a history of
533 coevolution with many interacting species, and may experience demographic bottlenecks or
534 genomic admixture of distant ~~sources~~ source populations. Thus, adaptive and random
535 evolutionary processes are likely intensified in introduced populations, and our understanding
536 of alien plants is incomplete without studying their genetics and evolution. Although this has
537 long been recognised (Baker & Stebbins 1965, Brown & Marshall 1981), genetic and
538 evolutionary studies of alien plants have increased only relatively recently (Bossdorf et al.
539 2005, Colautti & Lau 2015).

540

541 **5.1. Phenotypic Evolution of Alien Plants**

542 If introduced populations experience novel conditions, these will exert selection pressures on
543 plant phenotypes that may result in rapid evolution, provided there is genetic variation.

544 Identifying phenotypic divergence between native and introduced populations requires
545 approaches where offspring from different origins are grown in uniform environments, so
546 that a genetic basis to phenotypic differences can be confirmed. In recent decades, there has
547 been an explosion of common-garden studies with alien plants (reviewed in Bosssdorf et al.
548 2005, Colautti et al. 2009, Felker-Quinn et al. 2013). These studies most often test the
549 evolution of increased competitive ability (EICA) hypothesis, which proposes that some alien
550 plants may have become invasive because of evolutionary shifts from of reduced resource
551 allocation to defenses and increased allocation to growth and reproduction (Blossey &
552 Nötzold 1995). Some general patterns emerging from these studies are that (1) few studies
553 found results that are completely in line with the predictions of full support for EICA remains
554 rare (Felker-Quinn et al. 2013), but (2) significant differentiation often occurs in some growth
555 or defense traits between native and introduced plant populations (Bosssdorf et al. 2005,
556 Colautti et al. 2009, Felker-Quinn et al. 2013, Colautti & Lau 2015). Thus, rapid phenotypic
557 evolution appears to be common in introduced plant populations after introductions of plant
558 species into a new range.

559 Given the importance of herbivores and pathogens for plant evolution (Strauss &
560 Zangerl 2002) and the often observed release from specialist enemies (Mitchell & Power
561 2003, Meijer et al. 2016), shifts in allocation of resources from specialist to generalist enemy
562 defenses have been predicted in alien plants (shifting-defense hypothesis; Müller-Schärer et
563 al. 2004). Feeding experiments and chemical analyses of common-garden plants confirmed
564 that there are often differences in plant defenses between native and introduced populations,
565 and that defenses is-are reduced against specialists but not generalists (Doorduyn & Vrieling
566 2011, Felker-Quinn et al. 2013). Full support for the shifting-defense hypothesis has been
567 found in *Senecio jacobaea* where resistance to specialists is decreased but levels of
568 pyrrolizidine alkaloids and defense against generalists are increased in introduced populations

569 (Joshi & Vrieling 2005). In most cases the results are more complex, with evolutionary
570 changes in some aspects of defense but not others, shifts from constitutive to induced
571 defenses, or *vice versa*, and often large geographic variation within introduced and native
572 ranges (Orians & Ward 2010, Doorduyn & Vrieling 2011, Felker-Quinn et al. 2013).
573 However, when looking at the actual damage incurred by plants, a recent meta-analysis
574 showed that while resistance against specialist herbivores decreased in introduced
575 populations, resistance against generalist herbivores increased (Zhang et al. 2018). Taken
576 together, the evidence indicates that defense traits are key targets of selection and are
577 evolving rapidly in alien plants. This is also supported by experiments showing that exclusion
578 of herbivores can indeed lead to strong evolutionary responses within few generations (e.g.,
579 Agrawal et al. 2012).

Commented [WD2]: Do we want to use 'introduced' or 'alien' throughout?

580 ~~The common garden approach has limitations that were sometimes overlooked in~~
581 ~~earlier studies. Many of the early studies on evolutionary change in invasive plants suffer~~
582 ~~from several weaknesses.~~ Comparisons often involved low numbers of populations (Bossdorf
583 et al. 2005), and some comparisons were less meaningful as they confounded introduced and
584 native ranges with latitude or climate (Colautti et al. 2009). Comparisons were also often
585 made without knowing the sources of introduction, which can be misleading particularly for
586 species with one or few introductions. These pitfalls are now more widely recognized and
587 accounted for. Recent studies have used molecular data to identify sources of introductions
588 and then measured the phenotypic divergence between the source(s) and descendant
589 populations (e.g., Liao et al. 2014).

590 Simple common-garden studies can provide evidence for evolution but not adaptation,
591 and whether trait changes are really adaptive remains an open question (Figure 1). The latter
592 requires reciprocal transplant experiments or the combination of experiments with selection
593 measurements in wild populations. Both approaches are so far rare in the study of alien plants

Formatted: Font: Bold

594 (Colautti & Lau 2015). A recent analysis of published selection coefficients suggests that on
595 average, selection differentials are stronger in introduced than native populations, though data
596 for introduced species were few (Colautti & Lau 2015). Moreover, very few studies tested for
597 adaptive differentiation between ranges by transplanting plants between their native and
598 introduced ranges, and the few that did (e.g., Maron et al. 2004, Williams et al. 2008) found
599 little clear-cut evidence. All of these studies worked with only one or few sites per range,
600 limiting their power to draw general conclusions.

601 More studies (reviewed in Colautti & Lau 2015, [Oduor et al. 2016](#)) reciprocally
602 transplanted plants within the introduced range. Out of these studies, only one (Colautti &
603 Barrett 2013) on *Lythrum salicaria* found local adaptation in all study sites, whereas others
604 found ~~no or~~ only partial ~~or no~~ support for local adaptation. Nevertheless, [a meta-analysis of](#)
605 [such reciprocal transplant experiments within the introduced range of invasive species and](#)
606 [within the native range of native species](#) ~~the available evidence~~ indicates that adaptive
607 evolutionary processes are [at least as](#) common and as strong ~~in in introduced plant~~
608 ~~populations as in native ones~~ [invasive as in native species](#) (Oduor et al. 2016). [This is](#)
609 [surprising given that populations of invasive species had less time to adapt than populations](#)
610 [of native species, and supports the idea of an increased rate of evolutionary change in](#)
611 [invasive plants.](#)

612

613 5.2. Evolutionary Impacts on Native Species

614 Alien plants can also cause evolutionary responses in native residents. In particular, invasive
615 species are expected to exert selection on native species and cause evolutionary changes in
616 invaded communities (Strauss et al. 2006). While evolutionary studies on alien plants initially
617 focused entirely on alien evolution, recently attention has shifted toward evolutionary
618 responses of native species. For instance, native plants growing together with spotted

619 knapweed (*Centaurea maculosa*) in North America have evolved greater resistance to
620 knapweed allelopathy (Callaway et al. 2005). A combination of selection analyses and
621 reciprocal transplants in California showed that the exotic legume *Medicago polymorpha*
622 alters selection and adaptation of the native *Lotus wrangelianus* (Lau 2006, 2008). Perhaps
623 the most intriguing example of rapid evolution of a native species in response to an invasive
624 plant is the coevolutionary dynamic between invasive *Alliaria petiolata* and native
625 competitors in North America. It was shown that the [level of the](#) glucosinolate sinigrin
626 rapidly evolves in invasive *Alliaria* populations in response to native competitors, but the
627 native *Pilea pumila* equally rapidly evolves greater resistance to ~~these~~ [this](#) allelochemicals
628 (Lankau 2012). This could explain why glucosinolate concentrations decline with population
629 age in *Alliaria* populations in North America (Lankau et al. 2009). Rapid evolution [in](#)
630 [response](#) to invasive plants is common in native plants, as shown by a recent meta-analysis
631 (Oduor 2013). Broadening evolutionary studies of alien plants to their native competitors and
632 antagonists, and the coevolution between aliens and natives, is a promising and important
633 area for future research [\(Figure 1\)](#).

Formatted: Font: Bold

635 5.3. Molecular Genetics of Invasions

636 In addition to traits, the [ecology and](#) evolution of alien plants has also been studied through
637 molecular genetic analyses. Initially, these were mainly used for identifying pathways and
638 numbers of introductions, and genetic bottlenecks (reviewed in Bossdorf et al. 2005,
639 Dlugosch & Parker 2008). General insights from these studies are (1) contrary to
640 expectations, genetic diversity is often only moderately reduced in introduced populations,
641 (2) multiple introductions are common, and (3) admixture between different introductions can
642 even increase genetic diversity [in the introduced range](#). Since these earlier studies were

643 usually done with neutral markers, they were often disconnected from trait-based studies.
644 More recently, however, researchers have combined molecular with trait data, yielding more
645 powerful evolutionary studies.

646 Molecular data can provide knowledge of introduction pathways, allowing ~~better~~
647 ~~matching of alien populations with native ones for comparison to choose selection of the most~~
648 ~~likely native source populations as comparators for the invasive populations~~ (e.g., Liao et al.
649 2014). One can also incorporate population structure and stochastic processes into trait
650 analyses (Keller & Taylor 2008), analyse trait evolution along an invasion chronosequence
651 (Barker et al. 2017), test for phenotypic consequences of population admixture (Keller &
652 Taylor 2010), or compare specific hybrids or cytotypes of alien species (Hovick & Whitney
653 2014, Parepa et al. 2014).

654 Combining molecular and trait data can also ~~uncover~~ help to answer questions about
655 the genetic basis of evolving traits, i.e., the genes or genomic regions associated with
656 phenotypic changes in alien plants (Figure 1). Recent advances in sequencing technologies
657 allow high-resolution genomic data generation for any alien plant species, which can then be
658 used to construct genetic maps for Quantitative Trait Loci (QTL) or genome-wide association
659 studies. For example, Whitney et al. (2015) used a single-nucleotide-polymorphism map to
660 identify QTLs underlying fitness variation in invasive sunflowers. Gould & Stinchcombe
661 (2017) used whole-genome sequencing to show that flowering-time variation is associated
662 with different genes in the introduced versus native range of *Arabidopsis thaliana*. The use of
663 high-resolution genomic methods in invasion biology should be increased from now on.

Formatted: Font: Bold

5.4. The Way Forward for Research on the Evolution and Genetics of Invasive Plants

Much progress has been made in studying alien plant genetics and evolution. Rapid trait evolution is common in aliens as well as their native neighbours, but support for the EICA hypothesis is limited. Common garden studies have limitations, and are unlikely to further our understanding of alien plant evolution on their own. Future evolutionary studies of alien plants should (1) study selection and adaptation in the native versus introduced range, (2) explore coevolution between alien and native species, and (3) make combinations of trait and molecular data the standard for more targeted and mechanistic evolution studies (Figure 1).

Formatted: Font: Bold

Most evolutionary studies of alien plants have been done on ~~contemporary individuals~~ individuals representing the present generation of the populations rather than past generations, and were based on DNA sequence or trait data. However, two new research areas beyond these limits deserve mentioning, as they have the potential to greatly increase our understanding of alien plant evolution. First, with several hundred million specimens worldwide, herbaria represent huge historical archives of alien plant ~~evolutions~~. DNA sequence variation of herbaria specimen can be analysed with standard methods (see Vandepitte et al. 2014) for an example with alien plants and with specific ancient DNA methods (Suchan et al. 2016). Future research should attempt to take more advantage of this treasure. Second, individuals of the same species can also differ epigenetically, and there is currently much speculation about the role of epigenetic processes in rapid adaptation (Richards et al. 2017). Recent studies on *Alternanthera philoxeroides* (Gao et al. 2010) and *Fallopia japonica* (Zhang et al. 2017) found that genetically identical populations in the introduced ranges are often epigenetically differentiated, and that epigenetic variation can be associated with environment or phenotype. The paucity of studies precludes general conclusions, but they suggest epigenetic variation may contribute to the invasion success of these species. This should motivate researchers to study this possibility more thoroughly in future.

690

691 6. CONCLUSIONS

692 In a time of rapid environmental and biotic change, understanding why certain alien species
693 can successfully naturalize and become invasive has become a major objective in ecology
694 and evolution. At the same time, because alien organisms experience novel ecological
695 contexts, they provide study systems that advance our ecological and evolutionary
696 understanding. The ecology and evolution of alien plants is a very broad field crossing
697 different spatial and temporal scales, and with a plethora of hypotheses. We have highlighted
698 some of the major conceptual and methodological advances, as well as empirical results, that
699 we argue moved the field forward in the last decades or will do so in the near future.

700 Although we did not discuss modelling approaches, there have also been major advances in
701 developing more mechanistic models that bear great promise in predicting invasion risks
702 (e.g., Carboni et al. 2018). We call for more transparency on which ‘invasive’ definition
703 people use, and for a dissection of the different dimensions of invasiveness **(Figure 2)**. Using
704 the right comparators and accounting for introduction history may result in more consistent
705 findings. By integrating modern theories on plant coexistence and applying phylogenetic and
706 functional-trait approaches, ~~by integrating mechanistic processes into modelling approaches,~~
707 by studying the interplay between different biotic interactions, and by combining molecular
708 with experimental evolutionary approaches, we will advance our understanding of alien
709 plants and improve predictions on which aliens might invade in the future.

710

Formatted: Font: Bold

711 **DISCLOSURE STATEMENT**

712 The authors are not aware of any affiliations, memberships, funding, or financial holdings
713 that might be perceived as affecting the objectivity of this review.

714

715 **ACKNOWLEDGMENTS**

716 We thank Mark Vellend for inviting us to write this review. MvK, OB and WD thank the
717 German Research Foundation (DFG; projects KL 1866/9-1 and 10-1, BO 3241/1-1 and 8-1,
718 DA 1502/1-1) for funding.

719

720 **LITERATURE CITED**

- 721 Agrawal AA, Hastings AP, Johnson MTJ, Maron JL, Salminen JP. 2012. Insect herbivores
722 drive real-time ecological and evolutionary change in plant populations. *Science*
723 338:113-16
- 724 Aikio S, Duncan RP, Hulme PE. 2010. Lag-phases in alien plant invasions: separating the
725 facts from the artefacts. *Oikos* 119:370-78
- 726 Alpert P, Bone E, Holzapfel C. 2000. Invasiveness, invasibility and the role of environmental
727 stress in the spread of non-native plants. *Perspect. Plant Ecol. Evol. Syst.* 3:52-66
- 728 Atwater DZ, Ervine C, Barney JN. 2018. Climatic niche shifts are common in introduced
729 plants. *Nat. Ecol. Evol.* 2:34-43
- 730 Baker HG. 1965. Characteristics and modes of origins of weeds. In *The genetics of colonizing*
731 *species*, ed. HG Baker, GL Stebbins. New York, USA: Academic Press
- 732 Baker HG, Stebbins GL. 1965. *The Genetics of Colonizing Species*. New York: Academic
733 Press

734 Barker BS, Andonian K, Swope SM, Luster DG, Dlugosch KM. 2017. Population genomic
735 analyses reveal a history of range expansion and trait evolution across the native and
736 invaded range of yellow starthistle (*Centaurea solstitialis*). *Mol. Ecol.* 26:1131-47

737 Binggeli P. 2000. *Time-lags between introduction, establishment and rapid spread of*
738 *introduced environmental weeds*. Presented at Proceedings of the Third International
739 Weed Science Congress; 2000 June 6-11; Foz do Iguassu, Brazil

740 Blossey B, Nötzold R. 1995. Evolution of increased competitive ability in invasive
741 nonindigenous plants - a hypothesis. *J. Ecol.* 83:887-89

742 [Borer ET, Harpole WS, Adler PB, Lind EM, Orrock JL, Seabloom, Smith MD. 2014. Finding](#)
743 [generality in ecology: a model for globally distributed experiments. *Methods Ecol.*](#)
744 [Evol. 5:65-73](#)

745 Bossdorf O, Auge H, Lafuma L, Rogers WE, Siemann E, Prati D. 2005. Phenotypic and
746 genetic differentiation between native and introduced plant populations. *Oecologia*
747 144:1-11

748 [Boulangeat I, Georges D, Thuiller W. 2014. FATE HD: a spatially and temporally explicit](#)
749 [integrated model for predicting vegetation structure and diversity at regional scale.](#)
750 [Global Change Biol. 20:2368-78](#)

751 Caley P, Groves RH, Barker R. 2008. Estimating the invasion success of introduced plants.
752 *Divers. Distrib.* 14:196-203

753 Callaway RM, Ridenour WM. 2004. Novel weapons: invasive success and the evolution of
754 increased competitive ability. *Front. Ecol. Environ.* 2:436-43

755 Callaway RM, Ridenour WM, Laboski T, Weir T, Vivanco JM. 2005. Natural selection for
756 resistance to the allelopathic effects of invasive plants. *J. Ecol.* 93:576-83

757 Cappuccino N, Arnason JT. 2006. Novel chemistry of invasive exotic plants. *Biol. Lett.*
758 2:189-93

759 [Carboni M, Guéguen M, Barros C, Georges D, Boulangeat I, et al. 2018. Simulating plant](#)
760 [invasion dynamics in mountain ecosystems under global change scenarios. *Global*](#)
761 [Change Biol. 24:e289-e302](#)
762

763 Catford JA, Jansson R, Nilsson C. 2009. Reducing redundancy in invasion ecology by
764 integrating hypotheses into a single theoretical framework. *Divers. Distrib.* 15:22-40

765 CBD. 2000. *Alien Species that Threaten Ecosystems, Habitats or Species.*,
766 UNEP/CBD/COP/5/8, Secretariat of the Convention on Biological Diversity, Nairobi,
767 Kenya

768 Chesson P. 2000. Mechanisms of maintenance of species diversity. *Annu. Rev. Ecol., Evol.*
769 *Syst.* 31:343-66

770 Chun YJ, van Kleunen M, Dawson W. 2010. The role of enemy release, tolerance and
771 resistance in plant invasions: linking damage to performance. *Ecol. Lett.* 13:937-46

772 Colautti RI, Barrett SCH. 2013. Rapid adaptation to climate facilitates range expansion of an
773 invasive plant. *Science* 342:364-66

774 Colautti RI, Grigorovich IA, MacIsaac HJ. 2006. Propagule pressure: a null model for
775 biological invasions. *Biol. Invasions* 8:1023-37

776 Colautti RI, Lau JA. 2015. Contemporary evolution during invasion: evidence for
777 differentiation, natural selection, and local adaptation. *Mol. Ecol.* 24:1999-2017

778 Colautti RI, Maron JL, Barrett SCH. 2009. Common garden comparisons of native and
779 introduced plant populations: latitudinal clines can obscure evolutionary inferences.
780 *Evol. Appl.* 2:187-99

781 Cordeiro NJ, Patrick DAG, Munisi B, Gupta V. 2004. Role of dispersal in the invasion of an
782 exotic tree in an East African submontane forest. *J. Trop. Ecol.* 20:449-57

783 Crooks JA. 2005. Lag times and exotic species: the ecology and management of biological
784 invasions in slow-motion. *Ecoscience* 12:316-29

785 Cummings JA, Parker IM, Gilbert GS. 2012. Allelopathy: a tool for weed management in
786 forest restoration. *Plant Ecol.* 213:1975-89

787 Daehler CC. 2009. Short lag times for invasive tropical plants: evidence from experimental
788 plantings in Hawai'i. *PLoS One* 4:e4462

789 Darwin C. 1859. *On the Origin of Species by Means of Natural Selection, or, the*
790 *Preservation of Favoured Races in the Struggle for Life*. London: Murray

791 Davidson AM, Jennions M, Nicotra AB. 2011. Do invasive species show higher phenotypic
792 plasticity than native species and, if so, is it adaptive? A meta-analysis. *Ecol. Lett.*
793 14:419-31

794 Davis MA, Grime JP, Thompson K. 2000. Fluctuating resources in plant communities: a
795 general theory of invasibility. *J. Ecol.* 88:528-34

796 Dawson W, Burslem D, Hulme PE. 2009. Factors explaining alien plant invasion success in a
797 tropical ecosystem differ at each stage of invasion. *J. Ecol.* 97:657-65

798 Dawson W, Fischer M, van Kleunen M. 2012a. Common and rare plant species respond
799 differently to fertilisation and competition, whether they are alien or native. *Ecol. Lett.*
800 15:873-80

801 Dawson W, Keser LH, Winter M, Pyšek P, Kartesz J, et al. 2013. Correlations between
802 global and regional measures of invasiveness vary with region size. *NeoBiota* 16:59-
803 80

804 Dawson W, Moser D, van Kleunen M, Kreft H, Pergl J, et al. 2017. Global hotspots and
805 correlates of alien species richness across taxonomic groups. *Nat. Ecol. Evol.* 1:0186

806 Dawson W, Rohr RP, van Kleunen M, Fischer M. 2012b. Alien plant species with a wider
807 global distribution are better able to capitalize on increased resource availability. *New*
808 *Phytol.* 194:859-67

809 Dawson W, Schrama M. 2016. Identifying the role of soil microbes in plant invasions. *J. Ecol.*
810 104:1211-18

811 Dehnen-Schmutz K, Touza J, Perrings C, Williamson M. 2007. A century of the ornamental
812 plant trade and its impact on invasion success. *Divers. Distrib.* 13:527-34

813 Dietz H, Edwards PJ. 2006. Recognition that causal processes change during plant invasion
814 helps explain conflicts in evidence. *Ecology* 87:1359-67

815 Diez JM, Dickie I, Edwards G, Hulme PE, Sullivan JJ, Duncan RP. 2010. Negative soil
816 feedbacks accumulate over time for non-native plant species. *Ecol. Lett.* 13:803-09

817 Dlugosch KM, Parker IM. 2008. Founding events in species invasions: genetic variation,
818 adaptive evolution, and the role of multiple introductions. *Mol. Ecol.* 17:431-49

819 Doorduyn LJ, Vrieling K. 2011. A review of the phytochemical support for the shifting
820 defence hypothesis. *Phytochem. Rev.* 10:99-106

821 ~~Dullinger S, Gatttringer A, Thuiller W, Moser D, Zimmermann NE, et al. 2012. Extinction~~
822 ~~debt of high mountain plants under twenty-first century climate change. *Nat. Clim.*~~
823 ~~*Change* 2:619-22~~

824 Essl F, Dullinger S, Rabitsch W, Hulme PE, Hulber K, et al. 2011. Socioeconomic legacy
825 yields an invasion debt. *Proc. Natl. Acad. Sci. USA* 108:203-07

826 Felker-Quinn E, Schweitzer JA, Bailey JK. 2013. Meta-analysis reveals evolution in invasive
827 plant species but little support for Evolution of Increased Competitive Ability (EICA).
828 *Ecol. Evol.* 3:739-51

829 Feng YH, Maurel N, Wang ZH, Ning L, Yu FH, van Kleunen M. 2016. Introduction history,
830 climatic suitability, native range size, species traits and their interactions explain

831 establishment of Chinese woody species in Europe. *Global Ecol. Biogeogr.* 25:1356-
832 66

833 Funk JL. 2013. The physiology of invasive plants in low-resource environments. *Conserv.*
834 *Physiol.* 1:cot026

835 Gallien L, Carboni M. 2016. The community ecology of invasive species: where are we and
836 what 's next? *Ecography* 40:335-52

837 [Gallien L, Münkemüller T, Albert CH, Boulangéat I, Thuiller W. 2010. Predicting potential](#)
838 [distributions of invasive species: where to go from here? *Divers. Distrib.* 16:331-42](#)

839 Good RDO. 1931. A theory of plant geography. *New Phytol.* 30:149-71

840 Gould BA, Stinchcombe JR. 2017. Population genomic scans suggest novel genes underlie
841 convergent flowering time evolution in the introduced range of *Arabidopsis thaliana*.
842 *Mol. Ecol.* 26:92-106

843 Guisan A, Petitpierre B, Broennimann O, Daehler C, Kueffer C. 2014. Unifying niche shift
844 studies: insights from biological invasions. *Trends Ecol. Evol.* 29:260-69

845 [Guisan A, Zimmermann NE. 2000. Predictive habitat distribution models in ecology. *Ecol.*](#)
846 [Model. 135:147-86](#)

847 Harvey JA, Bukovinszky T, van der Putten WH. 2010. Interactions between invasive plants
848 and insect herbivores: a plea for a multitrophic perspective. *Biol. Conserv.* 143:2251-
849 59

850 Hayward J, Horton TR, Pauchard A, Nuñez MA. 2015. A single ectomycorrhizal fungal
851 species can enable a *Pinus* invasion. *Ecology* 96:1438-44

852 [Higgins SI, Richardson DM. 2014. Invasive plants have broader physiological niches. *Proc.*](#)
853 [Natl. Acad. Sci. USA](#) 111:10610-14

854 Hovick SM, Whitney KD. 2014. Hybridisation is associated with increased fecundity and
 855 size in invasive taxa: meta-analytic support for the hybridisation-invasion hypothesis.
 856 *Ecol. Lett.* 17:1464-77

857 Hufbauer R, Torchin M. 2007. Integrating ecological and evolutionary theory of biological
 858 invasions. In *Biological invasions*, ed. W Nentwig, pp. 79-96. Berlin: Springer-Verlag

859 Hyndman RJ, Mesgaran MB, Cousens RD. 2015. Statistical issues with using herbarium data
 860 for the estimation of invasion lag-phases. *Biol. Invasions* 17:3371-81

861 Joshi J, Vrieling K. 2005. The enemy release and EICA hypothesis revisited: incorporating
 862 the fundamental difference between specialist and generalist herbivores. *Ecol. Lett.*
 863 8:704-14

864 Keane RM, Crawley MJ. 2002. Exotic plant invasions and the enemy release hypothesis.
 865 *Trends Ecol. Evol.* 17:164-70

866 Keller SR, Taylor DR. 2008. History, chance and adaptation during biological invasion:
 867 separating stochastic phenotypic evolution from response to selection. *Ecol. Lett.*
 868 11:852-66

869 Keller SR, Taylor DR. 2010. Genomic admixture increases fitness during a biological
 870 invasion. *J. Evol. Biol.* 23:1720-31

871 Kempel A, Chrobock T, Fischer M, Rohr RP, van Kleunen M. 2013. Determinants of plant
 872 establishment success in a multispecies introduction experiment with native and alien
 873 species. *Proc. Natl. Acad. Sci. USA* 110:12727-32

874 Kowarik I. 1995. Time lags in biological invasions with regard to the success and failure of
 875 alien species. In *Plant Invasions - General Aspects and Special Problems*, pp. 15-38.
 876 Amsterdam: SPB Academic Publishing

877 Lankau RA. 2012. Coevolution between invasive and native plants driven by chemical
 878 competition and soil biota. *Proc. Natl. Acad. Sci. USA* 109:11240-45

879 Lankau RA, Nuzzo V, Spyreas G, Davis AS. 2009. Evolutionary limits ameliorate the
 880 negative impact of an invasive plant. *Proc. Natl. Acad. Sci. USA* 106:15362-67
 881 Larkin DJ. 2012. Lengths and correlates of lag phases in upper-Midwest plant invasions. *Biol.*
 882 *Invasions* 14:827-38
 883 Lau JA. 2006. Evolutionary responses of native plants to novel community members.
 884 *Evolution* 60:56-63
 885 Lau JA. 2008. Beyond the ecological: Biological invasions alter natural selection on a native
 886 plant species. *Ecology* 89:1023-31
 887 Le Roux JJ, Hui C, Keet JH, Ellis AG. 2017. Co-introduction vs ecological fitting as
 888 pathways to the establishment of effective mutualisms during biological invasions.
 889 *New Phytol.* 215:1354-60
 890 Levine JM, Adler PB, Yelenik SG. 2004. A meta-analysis of biotic resistance to exotic plant
 891 invasions. *Ecol. Lett.* 7:975-89
 892 Liao ZY, Zheng YL, Lei YB, Feng YL. 2014. Evolutionary increases in defense during a
 893 biological invasion. *Oecologia* 174:1205-14
 894 Lind EM, Parker JD. 2010. Novel weapons testing: are invasive plants more chemically
 895 defended than native plants? *PLoS One* 5:e10429
 896 Liu Y, van Kleunen M. 2017. Responses of common and rare aliens and natives to nutrient
 897 availability and fluctuations. *J. Ecol.* 105:1111-22
 898 Lockwood JL, Cassey P, Blackburn T. 2005. The role of propagule pressure in explaining
 899 species invasions. *Trends Ecol. Evol.* 20:223-28
 900 Lockwood JL, Hoopes MF, Marchetti MP. 2007. *Invasion ecology*. Malden, USA: Blackwell
 901 Publishing
 902 MacDougall AS, Gilbert B, Levine JM. 2009. Plant invasions and the niche. *J. Ecol.* 97:609-
 903 15

904 Macel M, de Vos RCH, Jansen JJ, van der Putten WH, van Dam NM. 2014. Novel chemistry
 905 of invasive plants: exotic species have more unique metabolomic profiles than native
 906 congeners. *Ecol. Evol.* 4:2777-86
 907 Mangla S, Inderjit, Callaway RM. 2008. Exotic invasive plant accumulates native soil
 908 pathogens which inhibit native plants. *J. Ecol.* 96:58-67
 909 Maron JL, Klironomos J, Waller L, Callaway RM. 2014. Invasive plants escape from
 910 suppressive soil biota at regional scales. *J. Ecol.* 102:19-27
 911 Maron JL, Vila M, Bommarco R, Elmendorf S, Beardsley P. 2004. Rapid evolution of an
 912 invasive plant. *Ecol. Monogr.* 74:261-80
 913 Maurel N, Hanspach J, Kühn I, Pyšek P, van Kleunen M. 2016. Introduction bias affects
 914 relationships between the characteristics of ornamental alien plants and their
 915 naturalization success. *Global Ecol. Biogeogr.* 25:1500-09
 916 Mayer K, Haeuser E, Dawson W, Essl F, Kreft H, et al. 2017. Current and future local
 917 naturalization potential of ornamental species planted in public green spaces and
 918 private gardens. *Biol. Invasions* DOI 10.1007/s10530-017-1594-y
 919 Mayfield MM, Levine JM. 2010. Opposing effects of competitive exclusion on the
 920 phylogenetic structure of communities. *Ecol. Lett.* 13:1085-93
 921 McKinney CL. 2001. Effects of human population, area, and time on non-native plant and
 922 fish diversity in the United States. *Biol. Conserv.* 100:243-52
 923 Meijer K, Schilthuizen M, Beukeboom L, Smit C. 2016. A review and meta-analysis of the
 924 enemy release hypothesis in plant-herbivorous insect systems. *PeerJ* 4:e2560v1
 925 Menzel A, Hempel S, Klotz S, Moora M, Pyšek P, et al. 2017. Mycorrhizal status helps
 926 explain invasion success of alien plant species. *Ecology* 98:92-102
 927 Mitchell CE, Power AG. 2003. Release of invasive plants from fungal and viral pathogens.
 928 *Nature* 421:625-27

929 Müller-Schärer H, Schaffner U, Steinger T. 2004. Evolution in invasive plants: implications
930 for biological control. *Trends Ecol. Evol.* 19:417-22

931 Oduor AMO. 2013. Evolutionary responses of native plant species to invasive plants: a
932 review. *New Phytol.* 200:986-92

933 Oduor AMO, Leimu R, van Kleunen M. 2016. Invasive plant species are locally adapted just
934 as frequently and at least as strongly as native plant species. *J. Ecol.* 104:957-68

935 Ollerton J, Watts S, Connerty S, Lock J, Parker L, et al. 2012. Pollination ecology of the
936 invasive tree tobacco *Nicotiana glauca*: comparisons across native and non-native
937 ranges. *J. Pollinat. Ecol.* 9:85-95

938 Ollerton J, Winfree R, Tarrant S. 2011. How many flowering plants are pollinated by animals?
939 *Oikos* 120:321-26

940 Orians CM, Ward D. 2010. Evolution of plant defenses in nonindigenous environments. *Annu.*
941 *Rev. Entomol.* 55:439-59

942 ~~Pagel J, Schurr FM. 2012. Forecasting species ranges by statistical estimation of ecological~~
943 ~~niches and spatial population dynamics. *Global Ecol. Biogeogr.* 21:293-304~~

944 Panetta FD, Mitchell ND. 1991. Homocline analysis and the prediction of weediness. *Weed*
945 *Res.* 31:273-84

946 Parepa M, Fischer M, Bossdorf O. 2013. Environmental variability promotes plant invasion.
947 *Nat. Commun.* 4:1604

948 Parepa M, Fischer M, Krebs C, Bossdorf O. 2014. Hybridization increases invasive knotweed
949 success. *Evol. Appl.* 7:413-20

950 Parker JD, Torchin ME, Hufbauer RA, Lemoine NP, Alba C, et al. 2013. Do invasive species
951 perform better in their new ranges? *Ecology* 94:985-94

952 Prior KM, Saxena K, Frederickson ME. 2014. Seed handling behaviours of native and
 953 invasive seed-dispersing ants differentially influence seedling emergence in an
 954 introduced plant. *Ecol. Entomol.* 39:66-74

955 Pyšek P, Jarošík V, Chytrý M, Danihelka J, Kühn I, et al. 2011. Successful invaders co-opt
 956 pollinators of native flora and accumulate insect pollinators with increasing residence
 957 time. *Ecol. Monogr.* 81:277-93

958 Pyšek P, Pergl J, Essl F, Lenzner B, Dawson W, et al. 2017. Naturalized alien flora of the
 959 world: species diversity, taxonomic and phylogenetic patterns, geographic distribution
 960 and global hotspots of plant invasion. *Preslia* 89:203-74

961 Pyšek P, Richardson DM. 2007. Traits associated with invasiveness in alien plants: where do
 962 we stand? In *Biological invasions*, ed. W Nentwig, pp. 97-125. Berlin: Springer-
 963 Verlag

964 Pyšek P, Richardson DM, Rejmánek M, Webster GL, Williamson M, Kirschner J. 2004.
 965 Alien plants in checklists and floras: towards better communication between
 966 taxonomists and ecologists. *Taxon* 53:131-43

967 Rabinowitz D. 1981. Seven forms of rarity. In *The Biological Aspects of Rare Plant*
 968 *Conservation.*, ed. H Synge. Chichester: John Wiley & Sons

969 Razanajatovo M, Maurel N, Dawson W, Essl F, Kreft H, et al. 2016. Plants capable of selfing
 970 are more likely to become naturalized. *Nat. Commun.* 7:13313

971 Razanajatovo M, van Kleunen M. 2016. Non-invasive naturalized alien plants were not more
 972 pollen-limited than invasive aliens and natives in a common garden. *Funct. Ecol.*
 973 30:1511-20

974 Reinhart KO, Tytgat T, van der Putten WH, Clay K. 2010. Virulence of soil-borne pathogens
 975 and invasion by *Prunus serotina*. *New Phytol.* 186:484-95

976 Rejmánek M. 1996. A theory of seed plant invasiveness: The first sketch. *Biol. Conserv.*
 977 78:171-81
 978 Rejmánek M. 2000. Invasive plants: approaches and predictions. *Austral Ecol.* 25:497-506
 979 Rejmánek M, Richardson DM. 2013. Trees and shrubs as invasive alien species-2013 update
 980 of the global database. *Divers. Distrib.* 19:1093-94
 981 Richards CL, Alonso C, Becker C, Bossdorf O, Bucher E, et al. 2017. Ecological plant
 982 epigenetics: Evidence from model and non-model species, and the way forward. *Ecol.*
 983 *Lett.* 20:1576-90
 984 Richardson DM, Pyšek P. 2006. Plant invasions: merging the concepts of species
 985 invasiveness and community invasibility. *Prog. Phys. Geog.* 30:409-31
 986 Richardson DM, Pyšek P, Rejmánek M, Barbour MG, Panetta FD, West CJ. 2000.
 987 Naturalization and invasion of alien plants: concepts and definitions. *Divers. Distrib.*
 988 6:93-107
 989 Sax DF, Stachowicz JJ, Brown JH, Bruno JF, Dawson MN, et al. 2007. Ecological and
 990 evolutionary insights from species invasions. *Trends Ecol. Evol.* 22:465-71
 991 Seabloom EW, Borer ET, Buckley YM, Cleland EE, Davies KF, et al. 2015. Plant species'
 992 origin predicts dominance and response to nutrient enrichment and herbivores in
 993 global grasslands. *Nat. Commun.* 6:7710
 994 Seabloom EW, Williams JW, Slayback D, Stoms DM, Viers JH, Dobson AP. 2006. Human
 995 impacts, plant invasion, and imperiled, plant species in California. *Ecol. Appl.*
 996 16:1338-50
 997 Seebens H, Blackburn TM, Dyer EE, Genovesi P, Hulme PE, et al. 2017. No saturation in the
 998 accumulation of alien species worldwide. *Nat. Commun.* 8

999 Seebens H, Essl F, Dawson W, Fuentes N, Moser D, et al. 2015. Global trade will accelerate
 1000 plant invasions in emerging economies under climate change. *Global Change Biol.*
 1001 21:4128-40
 1002 Shea K, Possingham HP. 2000. Optimal release strategies for biological control agents: an
 1003 application of stochastic dynamic programming to population management. *J. Appl.*
 1004 *Ecol.* 37:77-86
 1005 Shelby N, Duncan RP, van der Putten WH, McGinn KJ, Weser C, Hulme PE. 2016. Plant
 1006 mutualisms with rhizosphere microbiota in introduced versus native ranges. *J. Ecol.*
 1007 104:1259-70
 1008 Shipley B. 2000. *Cause and Correlation in Biology. A User's Guide to Path Analysis,*
 1009 *Structural Equations and Causal Inference.* Cambridge: Cambridge University Press
 1010 Simberloff D. 2009. The role of propagule pressure in biological invasions. *Annu. Rev. Ecol.,*
 1011 *Evol. Syst.* 40:81-102
 1012 [Simberloff D, Von Holle B. 1999. Positive interactions of nonindigenous species: Invasional](#)
 1013 [meltdown? *Biol. Invasions* 1:21-32](#)
 1014 Stinson KA, Campbell SA, Powell JR, Wolfe BE, Callaway RM, et al. 2006. Invasive plant
 1015 suppresses the growth of native tree seedlings by disrupting belowground mutualisms.
 1016 *PLoS Biol.* 4:727-31
 1017 Stohlgren TJ, Barnett DT, Kartesz JT. 2003. The rich get richer: patterns of plant invasions in
 1018 the United States. *Front. Ecol. Environ.* 1:11-14
 1019 Stout JC, Parnell JAN, Arroyo J, Crowe TP. 2006. Pollination ecology and seed production
 1020 of *Rhododendron ponticum* in native and exotic habitats. *Biodivers. Conserv.* 15:755-
 1021 77
 1022 Stout JC, Tiedeken EJ. 2017. Direct interactions between invasive plants and native
 1023 pollinators: evidence, impacts and approaches. *Funct. Ecol.* 31:38-46

1024 Strauss SY, Lau JA, Carroll SP. 2006. Evolutionary responses of natives to introduced
 1025 species: what do introductions tell us about natural communities? *Ecol. Lett.* 9:354-71
 1026 Strauss SY, Zangerl AR. 2002. Plant-insect interactions in terrestrial ecosystems. In *Plant*
 1027 *Animal Interactions: an Evolutionary Approach*, ed. CM Herrera, O Pellmyr, pp. 77-
 1028 106. Oxford: Blackwell Publishing
 1029 ~~Stricker KB, Harmon PE, Goss EM, Clay K, Flory SL. 2016. Emergence and accumulation of~~
 1030 ~~novel pathogens suppress an invasive species. *Ecol. Lett.* 19:469-77~~
 1031 ~~Talluto MV, Boulangeat I, Ameztegui A, Aubin I, Berteaux D, et al. 2016. Cross-scale~~
 1032 ~~integration of knowledge for predicting species ranges: a metamodeling framework.~~
 1033 ~~*Global Ecol. Biogeogr.* 25:238-49~~
 1034 Thuiller W, Gallien L, Boulangeat I, de Bello F, Munkemuller T, et al. 2010. Resolving
 1035 Darwin's naturalization conundrum: a quest for evidence. *Divers. Distrib.* 16:461-75
 1036 Traveset A, Richardson DM. 2014. Mutualistic interactions and biological invasions. *Annu.*
 1037 *Rev. Ecol., Evol. Syst.* 45:89-113
 1038 van Kleunen M, Dawson W, Bosssdorf O, Fischer M. 2014. The more the merrier: multi-
 1039 species experiments in ecology. *Basic Appl. Ecol.* 15:1-9
 1040 van Kleunen M, Dawson W, Essl F, Pergl J, Winter M, et al. 2015a. Global exchange and
 1041 accumulation of non-native plants. *Nature* 525:100-03
 1042 van Kleunen M, Dawson W, Maurel N. 2015b. Characteristics of successful alien plants. *Mol.*
 1043 *Ecol.* 24:1954-68
 1044 van Kleunen M, Dawson W, Schlaepfer D, Jeschke JM, Fischer M. 2010a. Are invaders
 1045 different? A conceptual framework of comparative approaches for assessing
 1046 determinants of invasiveness. *Ecol. Lett.* 13:947-58
 1047 van Kleunen M, Weber E, Fischer M. 2010b. A meta-analysis of trait differences between
 1048 invasive and non-invasive plant species. *Ecol. Lett.* 13:235-45

1049 Vandepitte K, De Meyer T, Helsen K, Van Acker K, Roldan-Ruiz I, et al. 2014. Rapid
 1050 genetic adaptation precedes the spread of an exotic plant species. *Mol. Ecol.* 23:2157-
 1051 64
 1052 Vilà M, Bartomeus I, Dietzsch AC, Petanidou T, Steffan-Dewenter I, et al. 2009. Invasive
 1053 plant integration into native plant-pollinator networks across Europe. *Proc. R. Soc.*
 1054 *Lond., Ser. B: Biol. Sci.* 276:3887-93
 1055 Vitousek PM, Walker LR. 1989. Biological invasion by *Myrica Faya* in Hawai'i: plant
 1056 demography, nitrogen fixation, ecosystem effects. *Ecol. Monogr.* 59:247-65
 1057 Von Holle B, Simberloff D. 2005. Ecological resistance to biological invasion overwhelmed
 1058 by propagule pressure. *Ecology* 86:3212-18
 1059 Wang B, Qiu YL. 2006. Phylogenetic distribution and evolution of mycorrhizas in land plants.
 1060 *Mycorrhiza* 16:299-363
 1061 Wang XY, Shen DW, Jiao J, Xu NN, Yu S, et al. 2012. Genotypic diversity enhances
 1062 invasive ability of *Spartina alterniflora*. *Mol. Ecol.* 21:2542-51
 1063 White PS, Pickett ST. 1985. Natural disturbance and patch dynamics: an introduction. In *The*
 1064 *Ecology of Natural Disturbance and Patch Dynamics*, ed. ST Pickett, PS White.
 1065 London: Academic Press Inc.
 1066 Whitney KD, Broman KW, Kane NC, Hovick SM, Randell RA, Rieseberg LH. 2015.
 1067 Quantitative trait locus mapping identifies candidate alleles involved in adaptive
 1068 introgression and range expansion in a wild sunflower. *Mol. Ecol.* 24:2194-211
 1069 Williams JL, Auge H, Maron JL. 2008. Different gardens, different results: native and
 1070 introduced populations exhibit contrasting phenotypes across common gardens.
 1071 *Oecologia* 157:239-48
 1072 Zhang YY, Parepa M, Fischer M, Bossdorf O. 2017. Epigenetics of colonizing species? A
 1073 study of Japanese knotweed in Central Europe. In *Invasion Genetics: the Baker &*

1074 *Stebbins Legacy*, ed. SCH Barrett, RI Colautti, KM Dlugosch, LH Rieseberg. Oxford:
1075 John Wiley & Sons

1076 [Zhang Z, Pant X, Blumenthal D, van Kleunen M, Liu M, Li B. 2018. Contrasting effects of](#)
1077 [specialist and generalist herbivores on resistance evolution in invasive plants. *Ecology*](#)
1078 [DOI:10.1002/ecy.2155](#)

Formatted: German (Germany)
Formatted: Indent: Left: 0 cm, Hanging: 1.25 cm
Formatted: English (United Kingdom)

Formatted: Indent: Left: 0 cm, First line: 0 cm

1080 **Figure caption**

1081

1082 **Figure 1**

1083 Visual summary of research intensity, consistency of results and open questions in the study of alien
1084 plant ecology and evolution, including general patterns and drivers, biotic interactions, and the role of
1085 evolution and genetics. We use a four-level heat-colour scale for the different cells to indicate whether
1086 in our opinion, research intensity, consistency of results and open questions have low, moderate, high
1087 or very high values. With arrows, we indicate the trends (decreasing, continuing, increasing, rapidly
1088 increasing) in the rate of research on each topic. For each topic, we list our top question that needs to
1089 be answered to better understand the ecology and evolution of alien plants.

1090

1091 **Figure 2**

1092 The stages and barriers (red bars) of the invasion process (after Richardson et al. 2000) and the
1093 multiple dimensions of invasiveness (e.g. Catford et al. 2016). When a species has been introduced
1094 from its native range across a biogeographic barrier it is an alien. Those aliens that passed the
1095 environmental barrier and are sometimes found in the wild but do not persist are casuals. The
1096 ones that passed the environmental and reproductive barriers and thus form persistent
1097 populations are naturalized species. A subset of the naturalized aliens is considered invasive.
1098 Invasiveness, however, is not a binary variable but a continuous one and has multiple dimensions (i.e.,
1099 distributional characteristics) indicated by the axes of the cube. Each of these invasiveness dimensions
1100 is associated with some main barrier along the continuum. Invasive species can theoretically occur
1101 anywhere within this multidimensional space. For example, species A is an invasive species that is a
1102 habitat specialist with a small range size that can become locally very abundant, due to overcoming
1103 competition from natives. Species B is an invasive species that has a large range through overcoming
1104 dispersal barriers, but is also a habitat specialist but has a large range and with a low local abundance.

Formatted: Line spacing: Double

Formatted: Line spacing: Double

1105 [Species C in an invasive species that has high values for each of the three invasiveness dimensions](#)
1106 [shown, and so has overcome competition, dispersal and environmental barriers.](#)